

A

NAME: _____ FAMILY PHYSICIAN: _____ AGE: _____ HT: _____ WT: _____

MEDICATIONS

SURGERY

MEDICAL HX

DO YOU SMOKE? Y/N
DO YOU DRINK ALCOHOL? Y/N
ARE YOU ALLERGIC TO LATEX? Y/N
DOES ANYONE IN YOUR FAMILY HAVE SKIN CANCER/MELANOMA? Y/N
IF SO, WHO?
WHAT MEDICATIONS ARE YOU ALLERGIC TO:

FEMALE PATIENTS:
LMP:
ARE YOU PREGNANT? Y/N
ARE YOU BREAST FEEDING? Y/N
WHEN WAS YOUR LAST MAMMOGRAM?

B

CC: SKIN GROWTH(S)/ACNE AS OUTLINED BELOW IN HPI BOX OR:

Table with 7 columns: LOCATION, PRESENT X, RECENT CHANGES, PREVIOUS TX, BLEEDING, PRURITIS/PAIN, GROWTH

PE: T P R BP
GEN: WD WN NAD
HEENT: NCAT EOMI
NECK: NT
CV: SR
PULM: CTA
ABD: SOFT NT ND
EXT: NO CCE
NEURO: MOTOR/SENS WNL
LAN: AX Y/N H/N Y/N ING Y/N

ROS:
WT LOSS N/V F/C
CHANGE IN VISION
SOB/DOE/CP
ABD PAIN/DIARRHEA
DYSURIA/HEMATURIA
SKIN RASH
DEPRESSION
WEAKNESS
BACK/JOINT PAIN

DERMATOLOGICAL FINDINGS AS OUTLINED IN BOX BELOW:

ACNE: PREVIOUS TX: CURRENT TX: PE:

Table with 12 columns: LOCATION, SIZE (CM), ELEVATED/FLAT, VERRUCOUS, COLOR, NODULAR, BLEEDING, PEDUNC., SMOOTH SCALLOPED, CYSTIC, TENDER, MOBILE

SKIN TAG BROWN/BLACK/TAN PEDUNCULATED NEOPLASMS AXILLA : (R) (L) NECK : (R) (L)
AK (S) RED/FLAT/FLAKY/OVAL SIZE CM LOCATION(S):
ABSCESS-TENDER FLUCTUANT MASS WITH OVERLYING ERYTHEMA LOCATED CM

NEOPLASM OF UNCERTAIN BEHAVIOR BCCA SCCA
SKIN TAGS SEBACEOUS CYST WART(S) MELANOMA
MALIGNANT NEOPLASM CUTANEOUS VASCULAR PROLIFERATION HYPERKERATOTIC LESION
ACNE HYPERHIDROSIS AXILLA ABSCESS HYPERTROPHIC/KELOID SCAR AK:

EXCISION LESION TODAY-SEE PROCED. NOTE EXCISION LESION SCHEDULED IN ASC
LIQUID NITROGEN LN was sprayed in a pulsed fashion onto the surface of the neoplasm(s) until the tissue turned white. The area(s) was(were) Treated, allowed to thaw, and re-treated for a total of 2 treatments to each lesion. Pt advised to f/u in 4 weeks if no resolution. Written post-procedure instructions.
BIOPSY: PARTIAL (P), SHAVE (S), SOFT TISSUE (ST) P/S/ST P/S/ST P/S/ST P/S/ST
TISSUE OBTAINED WITH#15 SCALPEL AFTER BETADINE PREP AND 1% LIDOCAINE INJECTION INTO AREA.PARTIAL BIOPSY OBTAINED A PORTION OF THE LESION, SHAVE BIOPSY REMOVED THE SURFACE OF THE LESION FLUSH WITH THE UNDERLYING TISSUE. SOFT TISSUE BIOPSY PERFORMED BY REMOVING AN ELIPSE OF THE LESION AND THE UNDERLYING SOFT TISSUE IF PE REVEALED A PALPABLE SOFT TISSUE COMPONENT.WOUND CLOSED WITH INTERRUPTED 4-0 CHROMIC.STERILE DRSG. WOUND CARE INSTR.
DESTRUCTION CUTANEOUS VASCULAR PROLIFERATION:BETADINE/1%LIDOCAINE.ELECTROCAUTERY APPLIED TO LESION RESULTING IN A BLACK DISCOLORATION OF THE AREA AND DISAPPEARANCE OF THE NEOPLASM. TOPICAL ABX OINT/STERILE DRSG. WRITTEN WOUND CARE INSTRUCTIONS PROVIDED.
I & D ABSCESS:BETADINE/1% LIDOCAINE. #11 SCALPEL USED TO INCISE ABSCESS. CC'S PURULENT FLUID.LOCULATIONS SEPARATED WITH MOSQUITO Clamp. Wound packed with sterile gauze. Wound care instructions provided. ANTIBIOTIC PRESCRIBED.
SKIN TAG(S):BETADINE PREP/1%LIDOCAINE.TAG(S) REMOVED WITH SCISSORS FLUSH WITH SURROUNDING EPIDERMIS. BASE CAUTERIZED.ABX OINT/DRSG.WOUND INSTRUC. Location(number): Axilla: R() L() Neck: R() L() INJECTION HYP/KELOID SCAR:40MG KENALOG/1CC 1%LIDOCAINE INJECTED
PARING OF BENIGN HYPERKERATOTIC LESION:BETADINE PREP.#15 SCALPEL USED TO PAR HYPERKERATOSIS FROM ITS BASE.TOPICAL ABX OINT/STERILE DRESG.
ACNE: medication(s) /skin care instructions-cleansing/moisturizing/med instructions.
MARC S. SCHEINER, M.D. DATE: F/U
ICD-9: 238.2 782.9 701.9 706.2 CPT: